

On consistency of determinants on cubic lattices

O. I. Mokhov

Consider the square lattice $\mathbb{Z}^2$ with vertices at points with integer-valued coordinates in $\mathbb{R}^2 = \{(x_1, x_2) | x_k \in \mathbb{R}, k = 1, 2\}$ and complex (or real) scalar fields u on the lattice $\mathbb{Z}^2$, $u : \mathbb{Z}^2 \rightarrow \mathbb{C}$, that are defined by their values $u_{i_1 i_2}$, $u_{i_1 i_2} \in \mathbb{C}$, at each vertex of the lattice with the coordinates (i_1, i_2) , $i_k \in \mathbb{Z}$, $k = 1, 2$. Consider a class of two-dimensional discrete equations on the lattice $\mathbb{Z}^2$ for the field u that are defined by functions $Q(x_1, x_2, x_3, x_4)$ of four variables with the help of the relations

$$Q(u_{ij}, u_{i+1,j}, u_{i,j+1}, u_{i+1,j+1}) = 0, \quad i, j \in \mathbb{Z}, \quad (1)$$

so that in each *elementary* 2×2 square of the lattice $\mathbb{Z}^2$, that is, in each set of vertices of the lattice with coordinates of the form (i, j) , $(i+1, j)$, $(i, j+1)$, $(i+1, j+1)$, $i, j \in \mathbb{Z}$, the value of the field u at one of vertices of the square is defined by the values of the field at three other vertices. In this case the field u on the lattice $\mathbb{Z}^2$ is completely determined by fixing initial data, for example, on the axes of coordinates of the lattice, u_{i0} and u_{0j} , $i, j \in \mathbb{Z}$. A particularly important role is played by *integrable nonlinear discrete equations*. In [1]–[3] integrable discrete equations of the form (1) are singled out by the very natural condition of *consistency on cubic lattices* (see also [4]–[9]). Consider the cubic lattice $\mathbb{Z}^3$ with vertices at points with integer-valued coordinates in $\mathbb{R}^3 = \{(x_1, x_2, x_3) | x_k \in \mathbb{R}, k = 1, 2, 3\}$ and fix initial data u_{i00} , u_{0j0} and u_{00k} , $i, j, k \in \mathbb{Z}$, on the axes of coordinates of the cubic lattice. A two-dimensional discrete equation (1) is called *consistent on the cubic lattice* if for initial data in general position the discrete equation (1) can be imposed in a consistent way on all two-dimensional sublattices of the cubic lattice $\mathbb{Z}^3$ at once (see [1]–[5]). Classifications of discrete equations of the form (1) that are consistent on the cubic lattice have been studied in [4] and [8] under some additional restrictions, see also [9]. The equation defined by determinants of 2×2 matrices of values of the field at vertices of elementary 2×2 squares of the lattice $\mathbb{Z}^2$:

$$u_{i,j+1}u_{i+1,j} - u_{i+1,j+1}u_{ij} = 0, \quad i, j \in \mathbb{Z}, \quad (2)$$

is an example of such two-dimensional nonlinear integrable discrete equation consistent on the cubic lattice. The equation (2) is linear with respect to each variable

¹The work was completed with the financial support of the Russian Foundation for Basic Research (grant no. 08-01-00054) and the Programme for Support of Leading Scientific Schools (grant no. NSh-1824.2008.1).

and invariant with respect to the full symmetry group of square. The fixing of arbitrary nonzero initial data u_{i0} and u_{0j} , $i, j \in \mathbb{Z}$, on the axes of coordinates of the lattice $\mathbb{Z}^2$ completely determines the field u on the lattice $\mathbb{Z}^2$ satisfying the discrete equation (2), and the fixing of arbitrary nonzero initial data u_{i00} , u_{0j0} and u_{00k} , $i, j, k \in \mathbb{Z}$, on the axes of coordinates of the lattice $\mathbb{Z}^3$ completely determines the field u on the lattice $\mathbb{Z}^3$ satisfying the discrete equation (2) on all two-dimensional sublattices of the cubic lattice $\mathbb{Z}^3$. The integrability (in the broad sense of the word) of the discrete equation (2) is obvious since it can be easily linearized: $\ln u_{i,j+1} + \ln u_{i+1,j} - \ln u_{i+1,j+1} - \ln u_{ij} = 0$, $i, j \in \mathbb{Z}$. In this work, we consider the question on consistency on cubic lattices for discrete nonlinear equations defined by determinants of matrices of higher orders (for orders $N > 2$). The condition of consistency on cubic lattices in the form as it was defined above is not satisfied for these discrete equations if $N > 2$. We prove that another, modified, condition of consistency on cubic lattices, which is proposed in this work, is satisfied for determinants of matrices of arbitrary orders.

Consider a discrete equation on the lattice $\mathbb{Z}^2$ defined by a relation for values of the field u at vertices of the lattice $\mathbb{Z}^2$ that form *elementary* 3×3 squares:

$$Q(u_{ij}, \dots, u_{i+s,j+r}, \dots, u_{i+2,j+2}) = 0, \quad 0 \leq s, r \leq 2, \quad i, j \in \mathbb{Z}. \quad (3)$$

The fixing of initial data u_{i0} , u_{i1} , u_{0j} and u_{1j} , $i, j \in \mathbb{Z}$, completely determines the field u on the lattice $\mathbb{Z}^2$ satisfying the discrete equation (3). Consider the cubic lattice $\mathbb{Z}^3$ and the condition of consistency on all two-dimensional sublattices of the cubic lattice $\mathbb{Z}^3$ for the discrete equation (3). Initial data can be specified, for example, at the following vertices of the lattice: u_{i00} , u_{i10} , u_{i01} , u_{i11} , u_{0j0} , u_{1j0} , u_{0j1} , u_{1j1} , u_{00k} , u_{10k} , u_{01k} and u_{11k} , $i, j, k \in \mathbb{Z}$. In the cube $\{(i, j, k), 0 \leq i, j, k \leq 2\}$, the values u_{202} , u_{212} , u_{220} , u_{221} , u_{022} and u_{122} are determined by the relations (3), and three relations must be satisfied for the value u_{222} on three faces of the cube at once. Consider the discrete nonlinear equation on the lattice $\mathbb{Z}^2$ defined by determinants of the matrices of values of the field u at vertices of the lattice $\mathbb{Z}^2$ that form elementary 3×3 squares:

$$\begin{aligned} & u_{i,j+2}u_{i+1,j+1}u_{i+2,j} + u_{i,j+1}u_{i+1,j}u_{i+2,j+2} + u_{i,j}u_{i+1,j+2}u_{i+2,j+1} - \\ & - u_{i,j}u_{i+1,j+1}u_{i+2,j+2} - u_{i,j+2}u_{i+1,j}u_{i+2,j+1} - u_{i,j+1}u_{i+1,j+2}u_{i+2,j} = 0, \end{aligned} \quad (4)$$

where $i, j \in \mathbb{Z}$. The equation (4) is linear with respect to each variable and invariant with respect to the full symmetry group of the configuration of vertices of the lattice $\mathbb{Z}^2$ that form elementary 3×3 squares. For initial data in general position, the discrete equation (4) is not consistent on two-dimensional sublattices of the cubic lattice $\mathbb{Z}^3$.

Consider another, modified, condition of consistency on cubic lattices for the discrete nonlinear equation (4). We shall require that the discrete equation (4) would be satisfied not only on all two-dimensional sublattices of the cubic lattice $\mathbb{Z}^3$, but

also on all unions of any two intersecting two-dimensional sublattices of the cubic lattice $\mathbb{Z}^3$, that is, the corresponding elementary 3×3 squares, on which the discrete equation (4) is considered, can be bent at right angle passing from one of two-dimensional sublattices to another, for example, $\{(i, 0, 0), (i, 1, 0), (i, 0, 1), i = 0, 1, 2\}$, $\{(0, j, 0), (1, j, 0), (0, j, 1), j = 0, 1, 2\}$ and $\{(0, 0, k), (1, 0, k), (0, 1, k), k = 0, 1, 2\}$. In this case, initial data can be specified, for example, at the following vertices of the cubic lattice $\mathbb{Z}^3$: $u_{i00}, u_{i01}, u_{2j0}, u_{2j1}, u_{10k}, u_{20k}, u_{110}$ and $u_{112}, i, j, k \in \mathbb{Z}$. We shall also call the corresponding discrete equations *consistent on cubic lattices*. Then the following theorem holds.

Theorem 1. *For arbitrary initial data in general position the nonlinear discrete equation (4) can be satisfied in a consistent way on all unions of any two two-dimensional sublattices of the cubic lattice $\mathbb{Z}^3$, that is, the discrete nonlinear equation (4) is consistent on the cubic lattice $\mathbb{Z}^3$.*

A similar property of consistency on cubic lattices holds for determinants of arbitrary orders N .

Theorem 2. *For any given integer N , the discrete nonlinear equation defined by determinants of the matrices of values of the field u at vertices of the lattice $\mathbb{Z}^2$ that form elementary $N \times N$ squares is consistent on the cubic lattice $\mathbb{Z}^3$.*

Bibliography

- [1] F.W. Nijhoff, A.J. Walker, *Glasgow Math. J.* **43A** (2001), 109–123.
- [2] F.W. Nijhoff, *Phys. Lett. A* **297** (2002), 49–58.
- [3] A.I. Bobenko, Yu.B. Suris, *Int. Math. Res. Notices* **11** (2002), 573–611.
- [4] V.E. Adler, A.I. Bobenko, Yu.B. Suris, *Comm. Math. Phys.* **233** (2003), 513–543.
- [5] A.I. Bobenko, Yu.B. Suris, arXiv: math/0504358.
- [6] A.I. Bobenko, Yu.B. Suris, *Uspekhi Mat. Nauk* **62:1** (2007), 3–50; English transl., A.I. Bobenko, Yu.B. Suris, *Russian Math. Surveys* **62:1** (2007), 1–43.
- [7] A.P. Veselov, *Uspekhi Mat. Nauk* **46:5** (1991), 3–45. English transl., A.P. Veselov, *Russian Math. Surveys* **46:5** (1991), 1–51.
- [8] V.E. Adler, A.I. Bobenko, Yu.B. Suris, arXiv: abs/0705.1663.
- [9] S.P. Tsarev, T. Wolf, *Lett. Math. Phys.* **84:1** (2008), 31–39.

Centre for Non-Linear Studies,
L.D.Landau Institute for Theoretical Physics,
Russian Academy of Sciences;
Department of Geometry and Topology,
Faculty of Mechanics and Mathematics,
M.V.Lomonosov Moscow State University;
E-mail: mokhov@mi.ras.ru; mokhov@landau.ac.ru; mokhov@bk.ru